

# Group meeting

## May 17<sup>th</sup>, 2024



# Instruction responsibilities

- Classes for Fall 2024:
  - PHY 331:
    - Need to make syllabus
  - PHY 361:
    - Need to make syllabus

# Service responsibilities

- Committee:
  - GlueX Compton Analysis Review Committee:
    - Waiting for author response

# Group responsibilities

- Undergrad: Met with Dylan on Tuesday
- Need to start writing DOE report that is due early June

# Analysis

TPOL:

- Waiting for new cooked files from 2022 data
- Need to run parallel TPOL analysis on 2022 data:
  - Standard
  - compressed

$KK\pi$ :

- Agave is alive but the Agave cluster is now dead ☹
- Need include polarization information in low-mass PWA



$E^*$ : Next slide

# $E^*$ Analysis

Cuts:

- Best combo



# $E^*$ $Q$ -factors



Fit with Voigtian:

- Center and width locked to PDG
- Resolution parameter  $\sigma$  allowed to vary

# $E^*$ $Q$ -factors




Fit with Voigtian:

- Center and width locked to PDG
- Resolution parameter  $\sigma$  allowed to vary

Found that the resolution was not consistent between 1<sup>st</sup> and 2<sup>nd</sup> Kaons


# $E^*$ $Q$ -factors



**Red** = first  $K^{+*}$

**Blue** = second  $K^{+*}$


# $E^*$ $Q$ -factors



**Red** = first  $K^{+*}$   
**Blue** = second  $K^{+*}$

- Red and blue consistent for fixed CL cut except at  $CL > 0.1$

# $E^*$ $Q$ -factors



**Red** = first  $K^{+*}$   
**Blue** = second  $K^{+*}$

- Red and blue consistent for fixed CL cut except at  $CL > 0.1$
- Generated  $Q$ -factors using resolution of **red** to cut out  $K^*$

# $E^*$ Analysis

Cuts:

- Best combo
- $CL > 0.1$

Table 1. Our estimate of the status of the  $\Xi$  resonances. Only those with an overall status of \*\*\* or \*\*\*\* are included in the Baryon Summary Table.

Particle	$J^P$	Overall status	Status as seen in —				
			$\Xi\pi$	$\Lambda K$	$\Sigma K$	$\Xi(1530)\pi$	Other channels
$\Xi(1318)$	1/2+	****					Decays weakly
$\Xi(1530)$	3/2+	****	****				
$\Xi(1620)$		**	**				
$\Xi(1690)$		***	**	***	**		
$\Xi(1820)$	3/2-	***	**	***	**	**	
$\Xi(1950)$		***	**	**		*	
$\Xi(2030)$		***		**	***		
$\Xi(2120)$		*		*			
$\Xi(2250)$		**					3-body decays
$\Xi(2370)$		**					3-body decays
$\Xi(2500)$		*		*	*		3-body decays

- \*\*\*\* Existence is certain, and properties are at least fairly well explored.
- \*\*\* Existence ranges from very likely to certain, but further confirmation is desirable and/or quantum numbers, branching fractions, *etc.* are not well determined.
- \*\* Evidence of existence is only fair.
- \* Evidence of existence is poor.


Table 1. Our estimate of the status of the  $\Xi$  resonances. Only those with an overall status of \*\*\* or \*\*\*\* are included in the Baryon Summary Table.

Particle	$J^P$	Overall status	Status as seen in —				
			$\Xi\pi$	$\Lambda K$	$\Sigma K$	$\Xi(1530)\pi$	Other channels
$\Xi(1318)$	$1/2+$	****					Decays weakly
$\Xi(1530)$	$3/2+$	****	****				
$\Xi(1620)$		**	**				
$\Xi(1690)$		***	**	***	**		
$\Xi(1820)$	$3/2-$	***	**	***	**	**	
$\Xi(1950)$		***	**	**		*	
$\Xi(2030)$		***		**	***		
$\Xi(2120)$		*		*			
$\Xi(2250)$		**					3-body decays
$\Xi(2370)$		**					3-body decays
$\Xi(2500)$		*		*	*		3-body decays

Looking in this range

- \*\*\*\* Existence is certain, and properties are at least fairly well explored.
- \*\*\* Existence ranges from very likely to certain, but further confirmation is desirable and/or quantum numbers, branching fractions, *etc.* are not well determined.
- \*\* Evidence of existence is only fair.
- \* Evidence of existence is poor.

# $\Xi^*$ Analysis




## $\Xi(1530)^0$ MASSES

### $\Xi(1530)^0$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1531.80 ± 0.32 OUR FIT</b>		Error includes scale factor of 1.3.		
<b>1531.78 ± 0.34 OUR AVERAGE</b>		Error includes scale factor of 1.4. See the ideogram below.		
1532.2 ± 0.7		DEBELLEFON 75B	HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi$
1533 ± 1		ROSS 73B	HBC	$K^- p \rightarrow \Xi \bar{K} \pi(\pi)$
1531.4 ± 0.8	59	BADIER 72	HBC	$K^- p$ 3.95 GeV/c
1532.0 ± 0.4	1262	BALTAY 72	HBC	$K^- p$ 1.75 GeV/c
1531.3 ± 0.6	324	BORENSTEIN 72	HBC	$K^- p$ 2.2 GeV/c
1532.3 ± 0.7	286	KIRSCH 72	HBC	$K^- p$ 2.87 GeV/c
1528.7 ± 1.1	76	LONDON 66	HBC	$K^- p$ 2.24 GeV/c
1532.1 ± 0.4	1244	ASTON 85B	LASS	$K^- p$ 11 GeV/c
1532.1 ± 0.6	2700	<sup>1</sup> BAUBILLIER 81B	HBC	$K^- p$ 8.25 GeV/c
1530 ± 1	450	BIAGI 81	SPEC	SPS hyperon beam

# $\Xi^*$ Analysis



$\Xi(1620)$

$I(J^P) = \frac{1}{2}(?)$  Status: \*  
 $J, P$  need confirmation.

OMITTED FROM SUMMARY TABLE


What little evidence there is consists of weak signals in the  $\Xi\pi$  channel. A number of other experiments (e.g., BORENSTEIN 72 and HASSALL 81) have looked for but not seen any effect.

## $\Xi(1620)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1620 OUR ESTIMATE</b>				
1624 ± 3	31	BRIEFEL	77	HBC $K^- p$ 2.87 GeV/c
1633 ± 12	34	DEBELLEFON	75B	HBC $K^- p \rightarrow \Xi^- \bar{K} \pi$
1606 ± 6	29	ROSS	72	HBC $K^- p$ 3.1-3.7 GeV/c



# $\Xi^*$ Analysis



## $\Xi(1690)$ MASSES

### MIXED CHARGES

VALUE (MeV)

DOCUMENT ID

**1690 ± 10 OUR ESTIMATE**

This is only an educated guess; the error given is larger than the error on the average of the published values.

### $\Xi(1690)^0$ MASS


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1686 ± 4	1400	ADAMOVICH 98	WA89	$\Sigma^-$ nucleus, 345 GeV/c
1699 ± 5	175	<sup>1</sup> DIONISI 78	HBC	$K^- p$ 4.2 GeV/c
1684 ± 5	183	<sup>2</sup> DIONISI 78	HBC	$K^- p$ 4.2 GeV/c

### $\Xi(1690)^-$ MASS



VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1691.1 ± 1.9 ± 2.0	104	BIAGI 87	SPEC	$\Xi^-$ Be 116 GeV
1700 ± 10	150	<sup>3</sup> BIAGI 81	SPEC	$\Xi^-$ H 100, 135 GeV
1694 ± 6	45	<sup>4</sup> DIONISI 78	HBC	$K^- p$ 4.2 GeV/c






# $E^*$ Analysis






# $E^*$ Analysis



# $\Xi^*$ Analysis



# $\Xi^*$ Analysis




- Bump structure becomes more pronounced after background subtraction

# $E^*$ Side Band check

- The side-band subtraction technique should yield similar results as the  $Q$ -factor method



# $\Xi^*$ Side Bands



**Blue** = Center of  $\Xi$  peak




**Red** = Side bands

# $\Xi^*$ Side Bands






**Blue** = Center of  $\Xi$  peak  
**Red** = Side bands

# $\Xi^*$ Side Bands



**Blue** = Center of  $\Xi^*$  peak  
**Red** = Side bands  
**Black** = Side band subtracted

# $\Xi^*$ Side Bands






**Blue** = Center of  $\Xi^*$  peak  
**Red** = Side bands  
**Black** = Side band subtracted

- Bump structure more pronounced after side-band subtraction



# $\Xi^*$ Side Bands



- Blue** = Center of  $\Xi^*$  peak
- Red** = Side bands
- Black** = Side band subtracted


- Bump structure more pronounced after side-band subtraction

**Need to run fitting routine over side-band subtracted results**



# $\Xi^*$ Analysis

- In progress



Sideband subtracted











