

# Isolation of the $\phi$ -mesons and $\Lambda(1520)$ from $e p \rightarrow e p K^+ K^-$ events

Patrick Walker


# Programs and procedures

- Data compiled from 112 runs

$$\phi \rightarrow K^+ K^-$$


- The peaks are not well defined.

Counts versus  $K^+ K^-$  mass




$$\Lambda(1520) \rightarrow p K^-$$


Counts versus  $p K^-$  mass





- Looking at the reaction:  
 $e p \rightarrow e p K^+ K^- X$ , where  $X$  is missing



- Cutting anything outside the red lines.




- After the cut from the missing mass. Individual particles can be distinguished from each other.
- Lambda 18XX stands for Lambda 1800 1810 1820 1830




# Before cut

Counts versus  $K^+ K^-$  mass




Counts versus  $p K^-$  mass




# After cut

Counts versus invariant  $K^+ K^-$  mass (cut)



Counts versus invariant  $p K^-$  mass (cut)



- We can now see distinguished peaks 😊



# Next steps

- Look to include
  - $K^0 \rightarrow \pi^+ \pi^-$
  - $\pi^0 \rightarrow \gamma \gamma$
  - neutrons
- Look for  $e p \rightarrow e p K^0 \Sigma^+$ , where  $\Sigma^+ \rightarrow p \pi^0$  or  $n \pi^+$
- Also interested in adding  $\pi^0 \rightarrow \gamma \gamma$